

h_c production at the Tevatron *

K. Sridhar[†]

*Theory Group, Tata Institute of Fundamental Research,
Homi Bhabha Road, Bombay 400 005, India.*

ABSTRACT

Non-Relativistic QCD (NRQCD) has provided a consistent framework for the physics of quarkonia. The colour-octet Fock components predicted by NRQCD have important implications for the phenomenology of quarkonium production in experiments. We have considered the consequences of the colour-octet channels for the production of the 1P_1 charmonium state, h_c , at large- p_T at the Tevatron. The colour-octet contributions to this state are found to be dominant and give a reasonably large rate for the production of h_c . This should make it feasible to look for this resonance in the $J/\psi + \pi$ decay channel. The observation of h_c is interesting in its own right, since its existence has yet to be experimentally confirmed. Moreover, this rate is a *prediction* of NRQCD, and the observation of the h_c at the Tevatron can, therefore, be used as a test of NRQCD.

*Presented at the XXXII Rencontres de Moriond, "QCD and High Energy Hadronic Interactions", March 22-29, 1997, Les Arcs, France.

[†]sridhar@theory.tifr.res.in

J/ψ production has traditionally been described in terms of the colour-singlet model [1, 2], which, in spite of its reasonable success with its predictions at lower energies, fails completely in explaining the data on large- p_T J/ψ production at the Tevatron [3]. This has led to a major revision of the theory of quarkonium formation. In the colour-singlet model, the relative velocity, v between the charm quarks is neglected. However, for charmonia $v^2 \sim 0.3$, and is not negligible. It is possible to include the effects coming from higher orders in a perturbation series in v in non-relativistic QCD (NRQCD) [4, 5].

In this picture, the physical charmonium state can be expanded in terms of its Fock-components in a perturbation series in v , and the $c\bar{c}$ states appear in either colour-singlet or colour-octet configurations in this series. The colour-octet $c\bar{c}$ state makes a transition into a physical state by the emission of one or more soft gluons, dominantly *via* $E1$ or $M1$ transitions with higher multipoles being suppressed by powers of v . The cross-section for the production of a meson H then takes on the following factorised form:

$$\sigma(H) = \sum_{n=\{\alpha,S,L,J\}} \frac{F_n}{m^{d_n-4}} \langle \mathcal{O}_\alpha^H(^{2S+1}L_J) \rangle \quad (1)$$

where F_n 's are the short-distance coefficients and $\mathcal{O}_n$ are local 4-fermion operators, of naive dimension d_n , describing the long-distance physics.

In fact, the importance of the colour-octet components was first noted [7] in the case of P -wave charmonium decays, and even in the production case the importance of these components was first seen [6] in the production of P -state charmonia at the Tevatron. The surprise was that even for the production of S -states such as the J/ψ or ψ' , where the colour singlet components give the leading contribution in v , the inclusion of sub-leading octet states was seen to be necessary for phenomenological reasons [8]. While the inclusion of the colour-octet components seem to be necessiated by the Tevatron charmonium data, the normalisation of these data cannot be predicted because the the long-distance matrix elements are not calculable. The data allow a linear combination of octet matrix-elements to be fixed [9, 10], and much effort has made recently to understand the implications of these colour-octet channels for J/ψ production in other experiments [11].

One important feature of the NRQCD Lagrangian is that it shows an approximate heavy-quark symmetry, which is valid to $O(v^2) \sim 0.3$. The implication of this symmetry is that the nonperturbative parameters have a weak dependence on the magnetic quantum number. We show that the production cross-section of the 1P_1 charmonium state, h_c , can be *predicted* in NRQCD, because of the heavy-quark symmetry. The Tevatron data on χ_c production fixes [10] the colour-octet matrix element which specifies the transition of a 3S_1 octet state into a 3P_J state. We would expect from heavy-quark spin symmetry of the NRQCD Lagrangian that the matrix-element for $^1S_0^{[8]} \rightarrow h_c$ should be of the same order as that for $^3S_1^{[8]} \rightarrow ^3P_1$. The production of the 1P_1 charmonium state, h_c , is, therefore, a crucial prediction of NRQCD.

The production of the h_c is interesting in its own right : charmonium spectroscopy [13] predicts this state to exist at the centre-of-gravity of the $\chi_c(^3P_J)$ states. While the E760 collaboration at the Fermilab has reported [14] the first observation of this resonance its existence needs further confirmation. In the following, we report on the calculations for h_c production at the Tevatron presented in Ref. [15].

Being a P -state, the leading colour-singlet contribution is already at $O(v^2)$, and at the same order we have the octet production of the 1P_1 state through an intermediate 1S_0 state. The octet channel is, therefore, expected to be very important for the production of this state. Once produced, the h_c can be detected by its decay into a $J/\psi + \pi^0$. The subprocesses that we are interested in are the following:

$$g + g \rightarrow ^1P_1^{[1]} + g,$$

$$\begin{aligned}
g + g &\rightarrow {}^1S_0^{[8]} + g, \\
q(\bar{q}) + g &\rightarrow {}^1S_0^{[8]} + q(\bar{q}), \\
q + \bar{q} &\rightarrow {}^1S_0^{[8]} + g.
\end{aligned}
\tag{2}$$

The ${}^1S_0^{[8]} \rightarrow h_c$ is mediated by a gluon emission in a $E1$ transition.

The cross-section $d\sigma/dp_T$ for h_c production at the Tevatron energy ($\sqrt{s} = 1.8$ TeV) presented in Ref. [15] are shown in Fig. 1. For 20 pb^{-1} total luminosity, for p_T integrated between 5 and 20 GeV we expect of the order of 650 events in the $J/\psi + \pi$ channel. Of these, the contribution from the colour-singlet channel is a little more than 40, while the octet channel gives more than 600 events. The colour-octet dominance is more pronounced at large- p_T . Recent results on J/ψ production from CDF are based on a total luminosity of 110 pb^{-1} . For this sample, more than 3000 events can be expected to come from the decay of the h_c into a J/ψ and a π . With this large event rate, the h_c should certainly be observable if the πi^0 coming from its decay can be reconstructed efficiently.

In conclusion, we find that a reasonably large rate for the production of the 1P_1 is expected at the Tevatron, with a dominant contribution from the colour-octet production channel. Heavy-quark symmetry relations allow us to infer the size of the non-perturbative matrix-elements and to predict the rate for h_c production and its subsequent decay into a J/ψ and a π . By looking at J/ψ events associated with a soft pion, it should be possible to pin down the elusive 1P_1 resonance at the Tevatron, given the large number of J/ψ events already available. We believe that this is a firm prediction of the NRQCD framework and may be a useful way of distinguishing this from other models of quarkonium formation.




Figure 1: $d\sigma/dp_T$ (in nb/GeV) for h_c production (after folding in with $\text{Br}(h_c \rightarrow J/\psi + \pi)=0.5$) at 1.8 TeV c.m. energy with $-0.6 \leq y \leq 0.6$.

References

- [1] E.L. Berger and D. Jones, *Phys. Rev.* **D 23** (1981) 1521.
- [2] R. Baier and R. Rückl, *Z. Phys.* **C 19** (1983) 251.
- [3] F. Abe et al., *Phys. Rev. Lett.* **69** (1992) 3704; *Phys. Rev. Lett.* **71** (1993) 2537; R. Demina, FERMILAB-CONF-96-201-E, Presented at the 11th Topical Workshop on Proton-Antiproton Collider Physics (PBARP 96), Padua, Italy, 26th May – 1st June 1996.
- [4] W.E. Caswell and G.P. Lepage, *Phys. Lett.*, **B 167** (1986) 437.
- [5] G.T. Bodwin, E. Braaten and G.P. Lepage, *Phys. Rev.* **D 51** (1995) 1125.
- [6] E. Braaten, M.A. Doncheski, S. Fleming and M. Mangano, *Phys. Lett.* **B 333** (1994) 548; D.P. Roy and K. Sridhar, *Phys. Lett.* **B 339** (1994) 141; M. Cacciari and M. Greco, *Phys. Rev. Lett.* **73** (1994) 1586.
- [7] G.T. Bodwin, E. Braaten and G.P. Lepage, *Phys. Rev.* **D 46** (1992) R1914.
- [8] E. Braaten and S. Fleming, *Phys. Rev. Lett.* **74** (1995) 3327.
- [9] M. Cacciari, M. Greco, M. Mangano and A. Petrelli, *Phys. Lett.* **B 356** (1995) 553.
- [10] P. Cho and A.K. Leibovich, *Phys. Rev.*, **D 53** (1996) 150; *Phys. Rev.*, **D 53** (1996) 6203.
- [11] N. Cacciari and M. Krämer, *Phys. Rev. Lett.* **76** (1996) 4128; J. Amundson, S. Fleming and I. Maksymyk, UTTG-10-95, hep-ph/9601298; S. Gupta and K. Sridhar, *Phys. Rev.* **D 54** (1996) 5545; M. Beneke and I. Rothstein, *Phys. Rev.* **D 54** (1996) 2005; W.-K. Tang and M. Vanttinen, NORDITA 96/18 P, hep-ph/9603266; S. Gupta and K. Sridhar, *Phys. Rev.* **D 55** (1997) 2650; E. Braaten and Y. Chen, *Phys. Rev. Lett.* **76** (1996) 730; K. Cheung, W.-Y. Keung and T.C. Yuan, *Phys. Rev. Lett.* **76** (1996) 877; P. Cho *Phys. Lett.* **B 368** (1996) 171; K. Cheung, W.-Y. Keung and T.C. Yuan, *Phys. Rev.* **D 54** (1996) 929; P. Ko, J. Lee and H.S. Song, *Phys. Rev.* **D 53** (1996) 1409; S. Fleming, O.F. Hernandez, I. Maksymyk and H. Nadeau, *Phys. Rev.* **D 55** (1997) 4098; K. Sridhar *Mod. Phys. Lett.* **A 11** (1996) 1555; see also M. Beneke, CERN Preprint CERN-TH/97-55, hep-ph/9703429 for a recent review and references therein.
- [12] P. Cho and M. Wise, *Phys. Lett.* **B 346** (1995) 129; M. Beneke and I. Rothstein, *Phys. Lett.* **B 372** (1996) 157; M. Beneke and M. Krämer, *Phys. Rev.* **D 55** (1997) 5269.
- [13] Y.-P. Kuang, S.F. Tuan and T.-M. Yan, *Phys. Rev.* **D 37** (1988) 1210.
- [14] T.A. Armstrong et al., *Phys. Rev. Lett.* **69** (1992) 2337.
- [15] K. Sridhar, *Phys. Rev. Lett.* **77** (1996) 4880.
- [16] R. Gastmans, W. Troost and T.T. Wu, *Nucl. Phys.* **B 291** (1987) 731.